Bleeding Control using a Combat Application Tourniquet

Task

Completed

Removed the C-A-T from the carrying pouch.	1 .		
Removed the C-A-T from the carrying pouch.	1st	2nd	3 rd
2 to the distriction of the dist	P / F	P/F	P / F
Slide the wounded extremity through the loop of the Self-Adhering Band or wrap around extremity.	P / F	P / F	P / F
Positioned the C-A-T above simulated wound site; left at least 2 inches of uninjured skin between the C-A-T and the wound site.	P / F	P/F	P / F
Twisted the Windlass Rod until the distal pulse was no longer palpable.	P/F	P/F	P / F
Locked the rod in place with the Windlass Clip.	P / F	P/F	P/F
Grasped the Windlass Strap, pulled it tight and adhered it to the Velcro on the Windlass Clip.	P/F	P/F	P / F
Verbalized using a marker to draw a "T" on the casualty's forehead and recorded the date and time the C-A-T was applied.	P/F	P/F	P / F
Did not secure the CAT properly for an arm/leg wound.			
lluator's Comment			
Imager's Comment			
	Date:		
ndent Name:	Date:		

Casualty Drags and Carries

Task		Complete	ed
Rescuer verbalizes for casualty to move to cover and provide self-aid	1st	2nd	3rd
Rescuer informs team and elicits assistance.	P / F	P / F	P/F
Rescuer assesses route and available cover	P / F	P / F	P/F
Determines appropriate carry.	P / F	P / F	P / F
Makes use of available cover and concealment during extraction	P / F	P/F	P/F
Does not cause further injury to the casualty	P / F	P/F	P / F

Evaluator's Comments:

Student Name:	Date:	
Evaluator:	Pass:	Fail:

Bleeding Control using a Combat Gauze Dressing

Completed

Task 2nd 3rd 1st P/F Verbalized: Do not use for minor wounds that do not entail severe P/F P / F Exposed the injury by cutting away the casualty's clothing. P/F P / F Opened the sterile Combat Gauze package. P/F P/F Placed the Combat Gauze directly into the wound where the bleeding was the heaviest. P/F P/F Held pressure on the bandage for 3 minutes. INSTRUCTOR: Inform the student that 3 minutes have passed. Reassess the wound to ensure that bleeding has stopped - leave the P/F P/F P/F Combat Gauze in place if bleeding is controlled. Verbalize that if there is continued bleeding, the Combat Gauze may be reconfigured in the wound or another Combat Gauze may be used in addition to the original. Applied a sterile pressure dressing over the bandage to secure it in P/FP/F P / F place. P / F P/F P/F Document and transport. Critical Criteria: Did not apply direct pressure for 3 minutes after application. Did not verbalize that failure to control hemorrhage requires that the Combat Gauze be reconfigured, or a second Combat Gauze used. Did not know that if Combat Gauze fails, the tourniquet re-tightened if the bleeding is on an extremity.

Student Name:	Date:	and the same of th
Evaluator:	Fai	1:

Airway Management Sucking Chest Wounds

Task				om	ւրյ	lete	ed		
		1		2	nc	đ		310	1
Verbalize: Seek hard cover to perform tactical field care.	P	1	F	P	1	F	P	1	F
Exposed the injury by cutting away the casualty's clothing.	P	1	F	P	/	F	P	1	F
Sweeps or whips excess blood from wound site.	P	1		4		F	P	1	F
Apply air tight dressing to wound.	P	1	F	P	1	F	P	/	F
INSTRUCTOR: Inform the student that patient was improved	breath	in	3				-		
Check for developing signs of difficulty in breathing.	l P	/	F				P		
Burps the dressing when confronted with signs of tension pneumothorax	P	/	F	P	1	F	P	/	F
Documented treatment.	P	1	F	P	1	F	P	1	F
Critical Criteria:									
Did not assess for exit wound.									
Did not apply occlusive dressing.									
Little Hot repair of the second secon									
Did not recognize developing tension pneumothorax.									
	ion.								
Did not recognize developing tension pneumothorax.	ion.								
Did not recognize developing tension pneumothorax. Failed to provide correct intervention for developing tens	ion.								
Did not recognize developing tension pneumothorax. Failed to provide correct intervention for developing tens	ion.								
Did not recognize developing tension pneumothorax. Failed to provide correct intervention for developing tens	ion.								
Did not recognize developing tension pneumothorax. Failed to provide correct intervention for developing tens	ion.								
Did not recognize developing tension pneumothorax. Failed to provide correct intervention for developing tens	ion.								

Evaluator:

Fail:

Pass:

Rapid Trauma Survey

		Complete	
	1 st	2nd	3rd
Verbalized: Maintaining situational awareness	P/F	P / F	P / I
Verbalized: Located in a position of hard cover.	P/F	P / F	P / I
Establishes patients level of consciousness using AVPU	P/F	P / F	P / I
Assess for active arterial bleeding	P/F	P / F	P / I
Applied tourniquet if indicated	P/F	P/F	P / I
Open the airway using head-tilt chin lift or jaw thrust maneuver	P / F	P / F	P / I
Assess patients breathing status.	P/F	P / F	P / F
Provide ventilations using pocket mask or mouth to mouth as indicated.	P / F	P/F	P / F
Correctly manage sucking chest wound as indicated.	P/F	P / F	P / F
Deploy wound packing for bleeding control if indicated	P/F	P / F	P / F
Preform a head to toe sweep to identify, locate and correct life threatening conditions.	P/F	P / F	P / F
Identifies Load and Go situations	P/F	P / F	P / F
Failure to accomplish any of the above. Failure to correctly address life threatening condition as soo	n a identi:	fied.	
Failure to accomplish any of the above. Failure to correctly address life threatening condition as soo	n a identi:	fied.	
* *	n a identi	fied.	
Failure to accomplish any of the above. Failure to correctly address life threatening condition as soo	n a identi	fied.	
Failure to accomplish any of the above. Failure to correctly address life threatening condition as soo	n a identi	fied.	
Failure to accomplish any of the above. Failure to correctly address life threatening condition as soo	n a identi	fied.	
Failure to accomplish any of the above. Failure to correctly address life threatening condition as soo	n a identi	fied.	
Failure to accomplish any of the above. Failure to correctly address life threatening condition as soo	n a identi:	fied.	

Airway Management CPR Mask/Barrier protection

		Complete	u
	1 st	2nd	3rd
Verbalized: Maintaining situational awareness	P / F	P / F	P / F
Verbalized: Located in a position of hard cover.	P/F	P / F	P / F
Establishes patients level of consciousness using AVPU	P/F	P / F	P / F
Open the airway using head-tilt chin lift or jaw thrust maneuver	P/F	P / F	P / F
Assess patients breathing status. LOOK, LISTEN, FEEL	P/F	P/F	P / F
Insert NPA as indicated.	P/F	P / F	
Provide ventilations using pocket mask or mouth to mouth as indicated.	P/F	P/F	P / F
Use EC grip in order to maintain good seal with mask.	P / F	P/F	P / F
Failure to accomplish any of the above.			
Failure to abide by NPA contraindacations.			
uator's Comments:			

Evaluator:

Fail:

Pass: